

On the Physical Interpretation of Malyshkin's (2008) Model of Resistive Hall-MHD Reconnection

Dmitri A. Uzdensky*

Princeton University/ Center for Magnetic Self-Organization (CMSO), Princeton, NJ 08544

(Dated: January 21, 2009)

A simple Sweet–Parker-like model for the electron current layer in resistive Hall magnetohydrodynamic (MHD) reconnection is presented, with the focus on the collisionless limit. The derivation readily recovers the main results obtained recently by Malyshkin [PRL, **101**, 225001 (2008)] and others, but is much quicker and more physically transparent. In particular, it highlights the role of resistive drag in determining the electron outflow velocity. The principal limitations of any such approach are discussed.

PACS numbers: 52.30.Cv, 52.30.Ex, 52.35.Vd, 94.30.cp

I. INTRODUCTION

Recently, Malyshkin [1] proposed an analytical model of reconnection in resistive Hall MHD regime without electron inertia (see also a recent paper by Simakov & Chaón 2008, Ref. [2]). Malyshkin's derivation was based on the local Taylor expansion of the resistive Hall MHD equations at the center of the electron layer. Under several simplifying assumptions, he was able to obtain expressions for the main parameters of the reconnection layer, including the reconnection rate E_z , in terms of a few input parameters, namely, the length of the resistive electron layer L , the magnetic field just outside the layer B_m , the density n , and the magnetic diffusivity $\eta = \eta' c^2 / 4\pi$ (where η' is the resistivity) [1]. One of the main results was that the reconnection rate becomes independent of the resistivity in the limit $\eta \rightarrow 0$ and scales inversely with L .

In this contribution we show that, under the assumptions of Malyshkin's model, his main results can be obtained much quicker and easier, in a way that we believe is more transparent and lends itself more readily to a clear physical interpretation. Specifically, we point out that Malyshkin's (2008) expressions for the reconnection layer parameters follow straight-forwardly from (essentially) a Sweet–Parker-like analysis applied to the inner electron layer. To make a direct comparison with Malyshkin's paper [1] easier, we adopt the same system of coordinates that he used, i.e., x is the direction across the layer, y is the outflow direction along the layer, and z is the ignorable direction. The only major modification in the Sweet–Parker procedure that one needs to make for the problem under consideration, is in the (electron) equation of motion (generalized Ohm's law) in the outflow (y) direction. Namely, since the electron mass is completely neglected here, the electron outflow velocity v_{ey} is no longer controlled by their inertia, as it would be in the classical Sweet–Parker approach. Instead, it is

determined by the balance between the outward acceleration due to the Lorentz force (and perhaps a comparable contribution from the pressure gradient force) and the collisional (resistive) drag exerted on the electrons by the much slower moving ions:

$$\frac{B_x j_{ez}}{c} \simeq -ne \eta' j_{ey} = n^2 e^2 \eta' v_{ey} \Rightarrow v_{ey} \sim \frac{B_x j_{ez}}{cn^2 e^2 \eta'}, \quad (1)$$

where the characteristic values for v_{ey} and B_x are taken at the outflow end of the electron layer, $x = 0, y = L$.

Next, using the z component of the generalized Ohm law (the electron equation of motion) both at the end of the electron layer ($x = 0, y = L$) and at the origin O ($x = 0 = y$), together with the stationarity condition $E_z(x, y) = \text{const}$, we can estimate $B_x \sim cE_z/v_{ey} = (c/v_{ey})\eta' j_{ez}$. Substituting this into equation (1), we immediately obtain

$$\begin{aligned} v_{ey} &\sim |v_{ez}| = j_{ez}/ne \sim cB_m/4\pi ne\delta \\ \Rightarrow v_{ey}\delta &\sim \frac{cB_m}{4\pi ne} = d_e V_{Ae} = d_i V_A, \end{aligned} \quad (2)$$

consistent with the expectation that the out-of-plane magnetic field $B_z \sim B_m$ (e.g., Refs. [3, 4]). Note that the resistivity has dropped out, because similar resistive terms enter in both the z and y components of the electron equation of motion, balancing the electric force $-neE_z$ and the Lorentz force $B_x j_{ez}/c$, respectively.

Combining equation (2) with the two remaining relationships of the standard Sweet–Parker analysis: the incompressibility condition, $|v_{ex}|L = v_{ey}\delta$, and the z -component of Ohm's law considered at the origin O and at the point $M = (x = \delta, y = 0)$ just above the resistive layer, $\eta B_m/\delta = cE_z = |v_{ex}|B_m \Rightarrow |v_{ex}|\delta = \eta$, we immediately get

$$\delta \sim \frac{\eta L}{d_i V_A} = \frac{L^2}{S d_i}, \quad (3)$$

$$|v_{ex}| \sim \frac{d_i}{L} V_A, \quad (4)$$

$$v_{ey} \sim |v_{ez}| \sim \frac{d_i V_A}{\delta} \sim V_A \frac{S d_i^2}{L^2}, \quad (5)$$

*Electronic address: uzdensky@astro.princeton.edu

$$c E_z \sim \frac{d_i}{L} V_A B_m, \quad (6)$$

which coincide with the corresponding Malyshkin's [1] expressions (20)-(26). Here, $S \equiv LV_A/\eta$ is the Lundquist number. Note that E_z is actually independent of the ion mass m_i . In addition, it is also independent of η , and the fundamental reason for this is that resistive drag controls the electron flow velocity in both the z and y directions in a similar way.

We note that recently similar results in resistive Hall-MHD were independently obtained by Simakov & Chacón [2], and even earlier by Wang et al. [5]. Furthermore, the expression (6) for the collisionless reconnection rate dates at least as far back as Ref. [6] (Malyshkin 2008, private communication).

Finally, we would like to stress that neither Malyshkin [1], nor Simakov & Chacón [2] address what, to us, seems to be the most important question in collisionless reconnection research: what determines the length L (in the outflow direction y) of the electron and ion layers? This question is similar to the old question about the length of the central diffusion region in the Petschek reconnection model. Both Malyshkin [1] and Simakov & Chacón [2] consider L (or w in Simakov & Chacón's notation) just an input parameter whose determination is beyond the scope of their papers. In our view, however, the question of whether the layer is macroscopic (independent of the local plasma parameters such as η , d_i , d_e , etc.) or microscopic, is of great importance, both fundamental and practical, since, for most space-

and astrophysical systems, the global system size is usually by many orders of magnitude larger than d_i , say. Thus, if L is comparable to the global system size, the resulting reconnection rate (6), even though independent of the resistivity, is too slow to explain many observable phenomena, such as solar flares. On the other hand, if L is microscopic (as seems to be indicated by numerical simulations [7, 8]; see, however, Refs. [5, 9]), it should be determined self-consistently as a part of the local analysis. Thus, until the issue of the length of the reconnection layer is resolved, achieving a final complete theory of collisionless reconnection cannot be claimed.

Likewise, neither the present analysis, nor the papers by Malyshkin [1] and Simakov & Chacón [2], attempt to estimate the magnetic field B_m just outside the electron layer in terms of the global reconnecting magnetic field B_0 measured just outside the ion layer (R. Kulsrud 2008, private communication). In general, depending on the strength of the electron and ion currents in the ion diffusion region ($\delta_e \ll x \ll \delta_i$), B_m may be much smaller than B_0 or comparable to B_0 . Until this issue is resolved, the solution of the reconnection problem remains incomplete.

Acknowledgments

I am very grateful to R. Kulsrud, M. Yamada, and L. Malyshkin for valuable discussions.

This work is supported by National Science Foundation Grant No. PHY-0215581 (PFC: Center for Magnetic Self-Organization in Laboratory and Astrophysical Plasmas).

-
- [1] L. M. Malyshkin, PRL, **101**, 225001 (2008)
 - [2] A. N. Simakov & L. Chacón, PRL, **101**, 105003 (2008)
 - [3] D. A. Uzdensky & R. Kulsrud, Phys. Plasmas, **13**, 062305 (2006)
 - [4] M. Yamada, Y. Ren, H. Ji, J. Breslau, S. Gerhardt, R. Kulsrud, A. Kuritsyn, Phys. Plasmas, **13**, 052119 (2006)
 - [5] X. Wang, A. Bhattacharjee, & Z. W. Ma, JGR, **105**, 27633 (2000)
 - [6] S. W. H. Cowley, in *Solar System Magnetic Fields*, edited by E. R. Priest (D. Reidel Publishing Company, 1985), 121.
 - [7] P. A. Cassak, M. A. Shay, & J. F. Drake, PRL, **95**, 235002 (2005)
 - [8] M. A. Shay, J. F. Drake, R. E. Denton, D. Biskamp, JGR, **103**, 9165 (1998)
 - [9] W. Daughton, J. Scudder, & H. Karimabadi, Phys. Plasmas, **13**, 072101 (2006)